

Journal of ORAL AND MAXILLOFACIAL SURGERY

IN THIS ISSUE

DENTOALVEOLAR SURGERY

No Difference in Surgical Outcomes Between Open and Closed Exposure of Palatally Displaced Maxillary Canines
Nicola A. Parkin, Chris Deery, et al

Effect of Different Methods for Decontaminating Tooth Enamel After Contact With Blood Before Bonding Orthodontic Buttons
Matheus Melo Pithon, Vanessa Oliveira Batista, et al

ANESTHESIA/FACIAL PAIN

Treatment Effectiveness of Arthrocentesis Plus Hyaluronic Acid Injections in Different Age Groups of Patients With Temporomandibular Joint Osteoarthritis
Luca Guarda-Nardini, Marco Olivo, et al

Mandibular Kinematics in Patients With Alloplastic Total Temporomandibular Joint Replacement—A Prospective Study
Sabine S. Linsen, Rudolf H. Reich, et al

DENTAL IMPLANTS

Zygomatic Bone-to-Implant Contact in 77 Patients With Partially or Completely Edentulous Maxillas
Thomas J. Balshi, Glenn J. Wolfinger, et al

Guided Bone Regeneration Using Cyanoacrylate-Combined Calcium Phosphate in a Dehiscence Defect: A Histologic Study in Dogs
Jung-Seok Lee, Seung-Hee Ko, et al

PATHOLOGY

Caldwell-Luc Operation Without Inferior Meatal Antrostomy: A Retrospective Study of 50 Cases
Yu-Chen Huang and Wen-Ho Chen

Cavernous Sinus Thrombosis: Current Therapy
Valmont Desa and Ryan Green

CRANIOMAXILLOFACIAL TRAUMA

The Financial Burden of Mandibular Trauma
Jasjit K. Dillon, Brian Christensen, et al

Does Traumatic Brain Injury Result in Accelerated Mandibular Fracture Healing?
Wei Huang, Zubing Li, et al

CRANIOMAXILLOFACIAL DEFORMITIES/COSMETIC SURGERY

Do Patients Treated With Bimaxillary Surgery Have More Stable Condylar Positions Than Those Who Have Undergone Single-Jaw Surgery?
Yoon-Ji Kim, Kyung-Min Oh, et al

Concomitant Removal of Mandibular Third Molars During Sagittal Split Osteotomy Minimizes Neurosensory Dysfunction
Jean-Charles Doucet, Archibald D. Morrison, et al

SURGICAL ONCOLOGY AND RECONSTRUCTION

Platelet-Rich Plasma Promotes Angiogenesis of Prefabricated Vascularized Bone Graft
Zhiwei Dong, Bei Li, et al

Long-Term Donor-Site Morbidity After Microsurgical Fibular Graft: Is There a Difference Between the Medial Approach and the Lateral Approach?
Philip Catalá-Lehnen, Carsten Rendenbach, et al

SPECIAL CONTRIBUTION

Summary of the Third Molar Clinical Trials: Report of the AAOMS Task Force for Third Molar Summary



Saving Faces | Changing Lives®

Official Journal of the
American Association of
Oral and Maxillofacial Surgeons

Zygomatic Bone-to-Implant Contact in 77 Patients With Partially or Completely Edentulous Maxillas

Thomas J. Balsbi, DDS, PhD,* Glenn J. Wolfinger, DMD,†
Nicolas J. Shuscavage, BA,‡ and Stephen F. Balsbi, MBE§

Purpose: Specifically with zygomatic implants, the bone-to-implant contact (BIC) at the zygomatic bone correlates with the survival of the implant because there is little anchorage at the alveolus. The purpose of this retrospective study was to view and measure the BIC of zygomatic implants in the zygomatic bone.

Materials and Methods: The patients in this study received zygomatic implants at a single private dental implant center. All patients were treated with the same immediate-loading protocol followed by postoperative cone beam computed tomography. The scans were exported to a computer-aided design system, where the BICs could be measured digitally. The BIC was analyzed by gender, and any statistical difference was determined by analysis of variance.

Results: The study sample was composed of 77 patients (62.3% women) receiving 173 zygomatic implants. The mean age of the sample was 59 ± 8.7 years. The mean BIC was 15.3 ± 5.6 mm (range, 4.9 to 32.9 mm) in the zygomatic bone. On average, $35.9\% \pm 11.7\%$ (range, 12.2% to 67.3%) of the implant came into contact with the zygomatic bone. The average BIC in men was 16.5 ± 6.0 mm, and the average BIC in women was 14.7 ± 5.4 mm, a statistically significant difference by analysis of variance ($P < .05$).

Conclusions: Evidence from this report indicates the zygomatic BIC varies greatly from patient to patient. These data show that the typical male patient has a greater zygomatic BIC than the typical female patient; however, these data do not support the hypothesis that the zygomatic BIC influences the zygomatic implant survival rate.

© 2012 American Association of Oral and Maxillofacial Surgeons
J Oral Maxillofac Surg 70:2065-2069, 2012

Anchorage is the primary concern when treating patients with dental and skeletal dysgnathia.¹ Patients presenting with complex anchorage problems, such as those with partially or completely edentulous jaws, achieve anchorage through osseointegration. Osseointegration implies a firm and lasting connection

between vital bone and titanium implants that is intended to distribute stress properly once connected to a permanent prosthesis.² Osseointegrated implants are proving to be a successful method for the replacement of lost teeth and related oral and facial structures.^{3,4}

Patients with severe atrophy of the maxilla present a complicated problem that cannot be treated with traditional dental implants alone. The techniques used in patients with little or no bone for implantation include the restoration of an atrophic maxilla by increasing the bone volume in necessary anatomic areas and improving the topography of existing bone. These results may be achieved by grafting bone from the patient's iliac crest,⁵ augmentation of the maxillary sinus,⁶ and Le Fort I osteotomies coupled with interpositional bone grafting.⁵ Although these techniques may have reasonable success rates, bone grafting procedures may be associated with intraoperative bleeding, postoperative infection, bone fracture,

Received from the PI Dental Center, Institute for Facial Esthetics, Fort Washington, PA.

*Founder and Prosthodontist.

†Prosthodontist.

‡Research Intern.

§Director of Research; President, CM Prosthetics, Inc, Fort Washington, PA.

Address correspondence and reprint requests to Mr S. F. Balsbi: 467 Pennsylvania Avenue, Suite 201, Fort Washington, PA 19034; e-mail: balsbi2@aol.com

© 2012 American Association of Oral and Maxillofacial Surgeons

0278-2391/12/7009-0\$36 00/0

<http://dx.doi.org/10.1016/j.joms.2012.05.016>

nerve dysfunction, perforation of the mucosa over the implant, the loss of a portion of the graft, ulcers, sinusitis, and wound dehiscence.^{7,8}

Recent surgical techniques have been adopted to shorten recovery time, decrease the cost of treatment, and drastically lessen surgical and recovery complications in all fields of medicine.^{7,9-15} One such technique involves the placement of implants in the zygomatic bone. Zygomatic implants provide craniofacial anchorage and allow for immediate prosthetic loading.¹⁶⁻²⁸

The stress caused by occlusal forces is supported mainly through the zygoma. The stress from these forces is transferred predominantly through the infra-zygomatic crest and divided to the frontal and temporal processes of the bone.^{28,29} Therefore, it is important to understand how much of the implant comes into contact with the zygomatic bone because the success of an immediately loaded implant is directly dependent on the primary stability provided by the bone and the eventual osseointegration of the titanium implant.

The purpose of this retrospective study was to examine the zygomatic bone-to-implant contact (BIC). This was examined because the authors hypothesized that the survival and osseointegration of these implants correlated directly to the anchorage necessary for immediate prosthetic delivery in patients with little to no maxillary bone.

In this study, the BIC was determined in a sample of patients receiving zygomatic implants and the factors associated with the BIC were investigated to determine the set of variables associated with the BIC. Some of these variables depend on the patient and others on the clinician. The specific aims of this study were to enroll a cohort of subjects receiving zygomatic implants, measure the BIC, and identify factors associated with the BIC.

Materials and Methods

STUDY DESIGN/SAMPLE

This retrospective study follows the guidelines for IRB exemption according to Ethical & Independent Review Services. To address the research purpose, the investigators designed and implemented a retrospective cohort study in which all patients who had zygomatic implants placed at a private clinical facility (PI Dental Center, Institute for Facial Esthetics, Fort Washington, PA) were analyzed. All patients were treated according to the Teeth in A Day protocol,^{30,31} that is, all implants were loaded within 2 hours of implant placement. Immediately after implant placement, the abutments were connected and an all-acrylic resin provisional prosthesis was delivered.³⁰

The exclusion criterion was limited to patients who underwent zygomatic implantation without postoperative cone beam computed tomographic examination. Cone beam computed tomographic scans (i-CAT, Imaging Sciences International, Hatfield, PA) were reviewed in Digital Imaging and Communications in Medicine (DICOM) format.

STUDY VARIABLE—GENDER

The range of BIC varies across patients because of the unique anatomic features of each patient. Because the BIC would differ in typical male and female patients from the obvious size differences, gender was the primary predictor variable in this study.

OUTCOME VARIABLE—BIC

The outcome variable in the present study was the BIC in the zygoma. In this retrospective cohort study, the BIC was determined using postoperative cone beam computed tomography and measuring the apical portion of the zygomatic implant in contact with the zygomatic bone. The export of the uncompressed DICOM data to the Procera computer-aided design system (NobelGuide; Nobel Biocare, Yorba Linda, CA) provided digital 3-dimensional images of each implant and the surrounding anatomic structures. The 3-dimensional images allowed digital manipulation of the computed tomographic images, making it possible to view the implants that had previously been placed in the zygoma. The Procera software then provided the images necessary to obtain the measurements of the zygomatic BIC. Once a clear image of the implant in the zygomatic bone was visible, measurements of the BIC were obtained by a digital manipulation of the 3-dimensional images (Fig 1A, B). Only the BIC in the zygoma was determined. All bone in the zygoma (cortical or trabecular) was considered in the calculation. The limited BIC in the maxilla was not analyzed.

DATA ANALYSES

Statistical analysis was performed using analysis of variance with a 95% confidence level ($P < .05$). The database was maintained in Excel (Microsoft, Redmond, WA), and analysis of variance was performed on the study variables using the add-on statistical package.

Results

Seventy-seven patients (31 men, 46 women; mean age, 59 ± 8.7 yr; age range, 33 to 80 yr) who underwent oral reconstruction because of severely atrophied maxillas were treated with 173 zygomatic implants (range, 1 to 4 implants). The overall implant survival rate was 96.5% (Table 1). Sixty-three of 66 implants (95.5%) placed in men survived, whereas




FIGURE 1. Screen captures from NobelGuide 3D Planning Software show zygomatic bone-to-implant contacts of A, 12.3 mm and B, 14.2 mm.

Balsbi et al. BONE-TO-IMPLANT CONTACT IN THE ZYGOMA. *J Oral Maxillofac Surg* 2012.

Table 1. LIFE TABLE ANALYSIS OF 173 IMMEDIATELY LOADED ZYGOMATIC IMPLANTS

Period	Implants	Failures	Survival Rate	Cumulative Survival Rate
0-3 mo	173	1	99.42%	99.42%
3-6 mo	173	3	98.27%	97.69%
6-9 mo	173	0	100.00%	97.69%
9-12 mo	173	1	99.42%	97.11%
1 yr	163	1	99.39%	96.53%
2 yr	144	0	100.00%	96.53%
3 yr	139	0	100.00%	96.53%
4 yr	87	0	100.00%	96.53%
5 yr	69	0	100.00%	96.53%
6 yr	55	0	100.00%	96.53%
7 yr	36	0	100.00%	96.53%
8 yr	28	0	100.00%	96.53%
9 yr	18	0	100.00%	96.53%
≥10 yr	16	0	100.00%	96.53%

Balsbi et al. BONE-TO-IMPLANT CONTACT IN THE ZYGOMA. *J Oral Maxillofac Surg* 2012.

104 of 107 implants (97.2%) placed in women survived. The survival rates between men and women were statistically similar ($P > .05$). The 6 zygomatic implant failures in this study are described in Table 2. All 3 zygomatic implant failures in men occurred in 1 patient, whereas 3 women each lost 1 zygomatic implant.

All completely edentulous patients were also treated with standard Brånemark System implants (Nobel Biocare) in the anterior maxilla. Some patients were also treated with Brånemark System implants in the pterygomaxillary region for added support and to eliminate distal cantilevers.^{32,33}

The zygomatic BIC (measured using the Procera computer-aided design/computer-assisted manufacturing software) showed a mean implant contact of 15.3 ± 5.6 mm (range, 4.9 to 32.9 mm) in all 77 patients. Zygomatic implants placed ranged from 30 to 52.5 mm. On average, $35.9\% \pm 11.7\%$ of the entire implant came into contact with the zygomatic bone. The BIC achieved is the basis of support for an imme-

Table 2. SIX ZYGOMATIC IMPLANT FAILURES

Gender	Zygomatic BIC (mm)	Implant in Zygoma (%)
F	16.3	40.75
F	17.1	38.0
F	12.2	28.7
M	14.4	36.0
M	18.3	34.8
M	17.8	33.9

Abbreviations: BIC, bone-to-implant contact; F, female; M, male.

Balsbi et al. BONE-TO-IMPLANT CONTACT IN THE ZYGOMA. *J Oral Maxillofac Surg* 2012.

diately loaded prosthesis. The percentage of the BIC ranged from 12.2% to 67.3%.

The average zygomatic BIC in the 29 men in this study was 16.5 ± 6.0 mm. The average zygomatic BIC in the 48 women in this study was 14.7 ± 5.4 mm. The 1.8-mm difference between men and women led to a 12.0% increase of the BIC in men. According to analysis of variance, there was a statistically significant difference in the BIC between men and women (95% confidence level; $P < .05$).

Discussion

The purpose of this retrospective study was to examine the BIC by measuring how much of the apical portion of the zygomatic implant was in contact with the zygoma. This appeared to be an important topic of study because with an increased understanding of the BIC comes an increased understanding of the factors related to osseointegration and successful immediate prosthetic loading. With an increase of zygomatic BIC, there is an overall inherent increase in prosthetic anchorage in patients with a resorbed or atrophic maxilla. It is the anchorage provided by the zygomatic implants that makes immediate prosthetic loading possible for these patients with severe atrophy. The advantages include the smaller number of surgeries (no grafting), shortened surgical and recovery times, and increased immediate prosthetic functionality compared with the alternative surgical and prosthetic options.

Patients with atrophy of the maxilla present a complicated clinical condition that cannot be treated with traditional dental implants alone. The lack of quality bone in the maxilla leaves little space for implants and does not ensure stability after prosthetic loading.

The zygomatic implants placed in this study had an overall cumulative survival rate (CSR) of 96.5% (167/173), proving a viable option for restoring the atrophic maxilla.¹⁷⁻²¹ The success of this technique resulted from the anchorage provided by the zygomatic implants (average BIC, 15.3 ± 5.6 mm). These data are comparable to the 2003 cadaver study performed by Van Steenberghe et al.³⁴ The present data indicated that as little as 4.9 mm of zygomatic BIC can immediately provide enough implant anchorage when the implant is rigidly connected to other implants in the dental arch. Permanent anchorage is achieved as osseointegration of the implants occurs. Therefore, implants placed in the zygoma obviate extensive and costly bone grafting procedures and provide superior prosthetic anchorage at the time of implant placement surgery.

The present data also indicated that BIC can differ by gender. In this study, male patients on average had 1.8 mm more BIC in the zygoma than did female patients. Although this result was statistically signifi-

cant, the increase in BIC in the male patients did not reflect a higher zygomatic implant cumulative survival rate (CSR).

Another variable that changes the zygomatic BIC is the angle at which the implant is placed. As the angle of the implant placement changes, the implant contacts different anatomic portions of the zygoma; this can lead to an increase or decrease of the BIC. The angle of implant insertion, which should be prosthetically driven, can be estimated by the examination of preoperative computed tomographic scans, but is subject to change because the implants are placed free hand by the surgeon. The angle of zygomatic implant placement was not a study variable because there is no anatomic reference point that is constant from patient to patient.

Although zygomatic implants usually are not the sole implants placed (although only 4 zygomatic implants have been used successfully to support a fixed prosthesis²²), these implants provide anchorage through an osseointegration that is predictable and superior to other implantation methods. This anchorage is necessary for immediate prosthetic support and is the backbone of an immediate-loading protocol for the severely atrophic maxilla. The high degree of fixed implant prosthetic success previously reported^{17,22,30} corroborates and validates the importance of the BIC measured and discussed in this report.

The evidence from this report indicated that the zygomatic BIC varies greatly from patient to patient. The present data showed that the typical male patient has a greater BIC than the typical female patient. Contrary to the primary study hypothesis, the limited sample size in this report showed that an increased zygomatic BIC did not result in increased cumulative survival rates of the zygomatic implants.

References

1. Jung BA, Yildizhan F, Wehrbein H: Bone-to-implant contact of orthodontic implants in humans—A histomorphometric investigation. *Eur J Orthod* 30:552, 2008
2. Adell R, Lekholm U, Rockler B, et al: 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 10:384, 1981
3. Koser LR, Campos PSF, Mendes CMC: Length determination of zygomatic implants using tridimensional computed tomography. *Braz J Oral Res* 20:4, 2006
4. Brånemark R, Brånemark PI, Rydevik B, et al: Osseointegration in skeletal reconstruction and rehabilitation: A review. *J Rehabil Res Dev* 38:175, 2001
5. Isaksson S, Alberius P: Maxillary alveolar ridge augmentation with onlay bone-grafts and immediate endosseous implants. *J Craniomaxillofac Surg* 20:2, 1992
6. Wood RM, Moore DL: Grafting of the maxillary sinus with intraorally harvested autogenous bone prior to implant placement. *Int J Oral Maxillofac Implants* 3:209, 1988
7. Higuchi KW: Minimization of bone grafting in the orofacial region, *in* Gottlander R, van Steenberghe D (eds): *Proceedings*

- of the First P-I Bränemark Scientific Symposium, Gothenburg 2009. Chicago, Quintessence, 2011, pp 125-135
8. Tolman DE: Reconstructive procedures with endosseous implants in grafted bone: A review of the literature. *Int J Oral Maxillofac Implants* 10:275, 1995
 9. Dasgupta P, Kirby RS: Outcomes of robotic assisted radical prostatectomy. *Int J Urol* 16:244, 2009
 10. Schell SR, Talamini MA, Udelsman R: Laparoscopic adrenalectomy for nonmalignant disease: Improved safety, morbidity, and cost-effectiveness. *Surg Endosc* 13:30, 1999
 11. Bencini L, Sanchez LJ, Boffi B, et al: Incisional hernia: Repair retrospective comparison of laparoscopic and open techniques. *Surg Endosc* 17:1546, 2003
 12. Briggs CD, Mann CD, Irving GR, et al: Systematic review of minimally invasive pancreatic resection. *J Gastrointest Surg* 13:1129, 2009
 13. Buchanan GN, Malik A, Parvaiz A, et al: Laparoscopic resection for colorectal cancer. *Br J Surg* 95:893, 2008
 14. Lotan Y, Cadeddu JA, Gettman MT: The new economics of radical prostatectomy: Cost comparison of open, laparoscopic and robot assisted techniques. *J Urol* 172:1431, 2004
 15. Poston RS, Tran R, Collins M, et al: Comparison of economic and patient outcomes with minimally invasive versus traditional off-pump coronary artery bypass grafting techniques. *Ann Surg* 248:638, 2008
 16. Branemark P-I, Breine U, Ardel R, et al: Intraosseous anchorage of dental prostheses. Part 1: Experimental studies. *Scand J Plast Reconstr Surg* 3:81, 1969
 17. Balshi SF, Wolfinger GJ, Balshi TJ: A retrospective analysis of 110 zygomatic implants in a single-stage immediate loading protocol. *Int J Oral Maxillofac Implants* 24:335, 2009
 18. Pi Urgell J, Revilla Gutiérrez V, Gay Escoda CG: Rehabilitation of atrophic maxilla: A review of 101 zygomatic implants. *Med Oral Patol Oral Cir Bucal* 13:E363, 2008
 19. Bedrossian E, Stumpel L III, Beckely ML, et al: The zygomatic implant: Preliminary data on treatment of severely resorbed maxillae. A clinical report. *Int J Oral Maxillofac Implants* 17:861, 2002
 20. Malevez C, Abarca M, Durdu F, et al: Clinical outcome of 103 consecutive zygomatic implants: A 6-48 months follow-up study. *Clin Oral Implants Res* 15:18, 2004
 21. Davo R, Malevez C, Rojas J, et al: Clinical outcome of 42 patients treated with 81 immediately loaded zygomatic implants: A 12-to-42 month retrospective study. *Eur J Implantol* 1:141, 2008
 22. Bedrossian E, Rangert B, Stumpel L, et al: Immediate function with the zygomatic implant: A graftless solution for the patient with mild to advanced atrophy of the maxilla. *Int J Oral Maxillofac Implants* 21:937, 2006
 23. Bothur S, Jonsson G, Sandahl L: Modified techniques using multiple zygomatic implants in reconstruction of the atrophic maxilla: A technical note. *Int J Oral Maxillofac Implants* 18:902, 2003
 24. Jensen OT, Adams MW: The maxillary M-4: A technical and biomechanical note for all-on-4 management of severe maxillary atrophy—Report of 3 cases. *J Oral Maxillofac Surg* 67:1739, 2009
 25. Aparacia C, Perales P, Rangert B, et al: Tilted implants as an alternative to maxillary sinus grafting: A clinical, radiologic and Periotest study. *Clin Implant Relat Res* 3:39, 2001
 26. Aparicio C, Ouazzani W, Hatano N: The use of zygomatic implants for prosthetic rehabilitation of the severely resorbed maxilla. *Periodontol* 2000 47:162, 2008
 27. Peñarocha M, García B, Martí E, et al: Rehabilitation of severely atrophic maxillae with fixed implant-supported prostheses using zygomatic implants placed using the sinus slot technique: Clinical report on a series of 21 patients. *Int J Oral Maxillofac Implants* 22:645, 2007
 28. Mattsson T, Köndell PA, Gynther GW, et al: Implant treatment without bone grafting in severely resorbed edentulous maxillae. *J Oral Maxillofac Surg* 57:281, 1999
 29. Ujjigawa K, Kato Y, Kizu Y, et al: Three-dimensional finite element analysis of zygomatic implants in craniofacial structures. *Int J Oral Maxillofac Surg* 36:620, 2007
 30. Balshi TJ, Wolfinger GJ: Teeth in a day immediate functional loading of dental implants. *Implant Dent* 10:231, 2002
 31. Sherry JS, Balshi TJ, Sims LO, et al: Treatment of a severely atrophic maxilla using an immediately loaded, implant-supported fixed prosthesis without the use of bone grafts: A clinical report. *J Prosthet Dent* 103:133, 2010
 32. Tulasne JF: Implant treatment of missing posterior dentition. *In* Albrektsson T, Zarb GA (eds): *The Bränemark Osseointegrated Implant*. Chicago, Quintessence, 1989, pp 103-115
 33. Balshi TJ, Wolfinger GJ, Balshi SF: Analysis of 356 pterygomaxillary implants in edentulous arches for fixed prosthesis anchorage. *Int J Oral Maxillofac Implants* 14:398, 1999
 34. Van Steenberghe D, Malevez C, Van Cleynenbreugel J, et al: Accuracy of drilling guides for transfer from three-dimensional CT-based planning to placement of zygoma implants in human cadavers. *Clin Oral Implants Res* 14:131, 2003